

Vampyrops lineatus. By Michael R. Willig and Robert R. Hollander

Published 27 February 1987 by The American Society of Mammalogists

Vampyrops lineatus (É. Geoffroy St.-Hilaire, 1810)

Geoffroy's Rayed Bat

Phyllostoma lineatum É. Geoffroy St.-Hilaire, 1810:180. Type locality restricted to Asunción, Paraguay, by Cabrera (1958).

Vampyrops lineatus, Peters, 1865:356. First use of current name combination.

Vampyrops nigellus Gardner and Carter, 1972:1. Type locality Huanhuachayo, Ayacucho, Perú.

CONTEXT AND CONTENT. Order Chiroptera, suborder Microchiroptera, superfamily Phyllostomoidea, family Phyllostomidae, subfamily Stenodermatinae. The genus *Vampyrops* contains eight (Jones and Carter, 1976, 1979) or nine (Honacki et al., 1982; Koopman, 1982; Swanepoel and Genoways, 1979) species; they may be distinguished on the basis of characters discussed in the diagnosis that follows. *Vampyrops lineatus* contains two recognized subspecies (Koopman, 1978):

V. l. lineatus (É. Geoffroy St.-Hilaire, 1810:180), see above (*sacrilus* Thomas is a synonym).

V. l. nigellus Gardner and Carter, 1972:1, see above.

DIAGNOSIS. *Vampyrops lineatus* can be distinguished morphometrically from most other congeners by both external and cranial features. It is larger than *V. helleri* and *V. brachycephalus* in seven cranial characters (greatest length of skull, condylobasal length, zygomatic breadth, postorbital constriction, breadth of braincase, length of maxillary toothrow, and breadth across upper molars) used by Swanepoel and Genoways (1979). Conversely, *V. lineatus* is smaller than *V. vittatus*, *V. infuscus*, and *V. aurarius* for each of the same seven characters. There is slight overlap between the largest specimens of *V. lineatus* and the smallest specimens of *V. dorsalis* in postorbital constriction and forearm length; *V. lineatus* is smaller than *V. dorsalis* in all other characters used. *Vampyrops umbratus* is considered conspecific with *V. dorsalis* (Jones and Carter, 1976; Rouk and Carter, 1972; Sanborn, 1955; Swanepoel and Genoways, 1979). Measurements of *V. recifinus* overlap those of *V. lineatus* in four (zygomatic breadth, breadth of braincase, length of maxillary toothrow, and breadth across upper molars) of seven cranial characters; as such, these taxa have been proposed as being either conspecific (Honacki et al., 1982; Rouk and Carter, 1972), or subspecific (Jones and Carter, 1976). Nevertheless, *V. lineatus* can be distinguished from *V. recifinus* by the lobation of the lower incisors (bilobed incisors in *V. lineatus* and trilobed incisors in *V. recifinus*; Carter and Dolan, 1978).

GENERAL CHARACTERS. In external appearance, *V. lineatus* is like other species of *Vampyrops* in resembling *Artibeus* somewhat (Fig. 1). In general, the muzzle is short, wide, and flattened; the noseleaf is well developed and lanceolate, the tip is pointed whereas the base resembles a horseshoe with a finely crenulated free margin. The rounded ears are about the size of the head; their internal margins are convex whereas the external margins are concave. The tragus is well developed, pointed, and crenulated along the internal margin. Wing membranes connect at the metacarpals, almost at the base of the phalanges. The calcar is short. The interfemoral membrane is strongly indented with the edge of the concavity touching at the knees. The thumb is long with a well-developed nail. The dorsal pelage is thick and long (5 to 6 mm). Sparse hairs cover the superior surface of the interfemoral membrane, the legs, and the digits of the feet. A fringe of fine hairs extends along the posterior margin of the interfemoral membrane. The digits of the feet are small with weak nails. The dorsum is yellow-brown to dark-brown whereas the venter is paler. The head has four broad white stripes (Fig. 1); two extend from the base of the noseleaf to

the posterior edge of the ear and two occur on the cheek below the eye. A distinct white stripe extends along the dorsum from the occiput to the base of the interfemoral membrane.

The dental formula is i 2/2, c 1/1, p 2/2, m 3/3, total 32. Upper incisors are large, in contact, and oblique, each with a broad cusp projecting inward and slightly forward (Fig. 2). The outer upper incisors are small, approximately half the size of I1, conical, and straight. The lower incisors are small, equal in size, and bifid. The canines and premolars are similar to those of *Artibeus*. The surface of the crowns of the lower molars is almost smooth; M3 is reduced. The cranium is similar to that of *Artibeus*, although the rostrum is relatively longer and narrower.

In addition to reporting measurements of *V. lineatus*, Swanepoel and Genoways (1979) listed a number of other studies that contain mensural data (Allen, 1891; Chuna Vieira, 1942; Dobson, 1878; Elliot, 1904; Goodwin, 1946; Hershkovitz, 1949; Lima, 1926; Sanborn, 1955). Taddei (1979) reported statistically significant secondary sexual variation in samples from northwestern São Paulo, Brazil, for 2 of 18 cranial characters (palate width and basal length) and 5 of 18 external characters (forearm length, length of the metacarpal of digit III, length of the first phalanx of digit III, length of the third phalanx of digit III, and length of the metacarpal of digit IV). Willig (1983) examined morphometric variation within and between populations in northeast Brazil; 9 of 14 external characters (total length, hind foot length, forearm length, weight, length of digit I, length of digit III, length of digit IV, length of digit V, tibia length) and none of 16 cranial characters exhibited significant secondary sexual variation. Significant differences between populations from Caatingas and edaphic Cerrado habitats were obtained for four cranial characters (rostral breadth, breadth across the upper molars, upper molariform toothrow length, width of widest molar) and four external characters (hind foot length, forearm length, length of digit IV, length of digit V). Mean values (followed by standard deviations in parentheses) in mm (Willig, 1983), each based upon a sample of 20 individuals, for Caatingas males, Caatingas females, Cerrado males, and Cerrado females, respectively, are: total length, 64.40 (2.98), 67.85 (2.60), 63.55 (2.78), 66.70 (2.66); hind foot length, 11.20 (0.62), 11.60 (0.60), 11.55 (0.51), 11.90 (0.45); ear length, 18.05 (1.05), 18.15 (0.67), 17.80 (0.89), 18.15 (0.67);




FIG. 1. Photograph of a live specimen of *Vampyrops l. lineatus* from northeast Brazil (photograph by M. R. Willig).



FIG. 2. Dorsal, ventral, and lateral views of cranium and lateral view of lower jaw of *Vampyrops l. lineatus* (M. R. Willig field number 1931 ♂), from O Aeroporto de Crato, Crato, Ceará. Greatest length of skull is 18.9 mm (photograph by N. Olson).

tragus length, 7.15 (0.75), 7.35 (0.59), 7.30 (0.57), 7.10 (0.72); forearm length, 47.20 (1.94), 47.45 (2.04), 46.00 (1.12), 47.20 (1.06); length of digit I, 10.65 (0.75), 10.95 (0.76), 10.60 (0.75), 11.10 (0.64); length of digit III, 96.80 (3.49), 97.10 (3.70), 94.00 (2.64), 97.35 (2.41); length of digit IV, 70.95 (2.04), 71.90 (2.55), 69.35 (2.03), 71.05 (1.93); length of digit V, 65.75 (1.94), 67.25 (2.86), 64.35 (1.90), 66.10 (1.65); tibia length, 18.00 (0.46), 18.35 (0.88), 17.95 (0.76), 18.50 (0.83); calcar length, 4.75 (0.85), 4.30 (0.57), 4.25 (0.44), 4.30 (0.47); noseleaf length, 6.95 (0.76), 7.00 (0.86), 6.85 (0.88), 7.25 (0.64); greatest length of skull, 24.32 (0.46), 24.40 (0.37), 24.38 (0.56), 24.41 (0.35); condylobasal length, 21.90 (0.44), 22.00 (0.46), 21.83 (0.61), 21.89 (0.38); zygomatic breadth, 14.32 (0.34), 14.37 (0.40), 14.25 (0.37), 14.17 (0.39); postorbital constriction, 6.26 (0.19), 6.29 (0.20), 6.23 (0.18), 6.22 (0.25); mastoid breadth, 12.12 (0.26), 12.22 (0.31), 12.21 (0.24), 12.16 (0.32); breadth of braincase, 10.51 (0.29), 10.56 (0.26), 10.56 (0.23), 10.58 (0.16); rostral breadth, 7.46 (0.26), 7.42 (0.31), 7.03 (0.36), 6.86 (0.21); height of braincase, 11.48 (0.23), 11.42 (0.31), 12.44 (0.29), 12.36




FIG. 3. Distribution of *Vampyrops lineatus* (principally after Koopman, 1983). Subspecies are *V. l. lineatus* (crosshatch) and *V. l. nigellus* (diagonal).

(0.24); breadth across upper molars, 10.19 (0.23), 10.40 (0.31), 10.17 (0.25), 10.13 (0.25); breadth across upper canines, 6.05 (0.22), 6.15 (0.23), 6.09 (0.15), 6.02 (0.21); length of maxillary tooththrow, 8.59 (0.33), 8.75 (0.32), 8.76 (0.28), 8.62 (0.27); length of upper molariform tooththrow, 7.09 (0.26), 7.16 (0.28), 7.05 (0.21), 7.98 (0.20); width of widest molar, 2.51 (0.11), 2.53 (0.13), 2.42 (0.14), 2.45 (0.16); greatest length of mandible, 16.62 (0.41), 16.74 (0.28), 16.49 (0.45), 16.49 (0.32); length of mandibular tooththrow, 9.53 (0.26), 9.60 (0.33), 9.54 (0.25), 9.46 (0.26); and length of coronoid process, 5.69 (0.25), 5.87 (0.25), 5.76 (0.29), 5.74 (0.21). Mean weights (g) followed by standard deviations in parentheses for the same series are 23.93 (1.32), 26.44 (3.73), 23.10 (1.78), 26.48 (2.90).

DISTRIBUTION. The genus *Vampyrops* occurs from subtropical South America north to southern Mexico (Fig. 3). The two recognized subspecies of *V. lineatus* are not parapatric (Anderson et al., 1982). *Vampyrops l. nigellus* occurs along the Andes from northern Colombia south through Ecuador and Peru to western Bolivia. *Vampyrops l. lineatus* occurs from northeastern Brazil across to central Bolivia and south through eastern Paraguay and northeastern Argentina to northern Uruguay. There is no known fossil record.

ONTOGENY AND REPRODUCTION. Although the family Phyllostomidae is characterized by a single young per litter, the presence of a developed corpus luteum in each ovary of nulliparous females of *V. lineatus* suggests the possibility of occasional twinning in this species (Taddei, 1976). Taddei (1973) obtained mensural data on a neonate within 24 h of parturition, the measurements (in mm, and mass in g) and percentage of maternal size, respectively, follow: head and body length 46.5, 68.3; ear length, 14.5, 74.3; tragus length, 4.5, 69.2; forearm length, 27.0, 58.0; length of digit I, 6.5, 86.6; length of digit III, 21.0, 48.8; tibia length, 12.0, 68.5; foot length, 10.5, 95.4; weight, 8.30, 30.8. Embryologic development has not been examined otherwise.

Peracchi and Albuquerque (1971) captured pregnant females

in December, January, and March in the Brazilian state of Rio de Janeiro. Taddei (1973, 1976) obtained pregnant females in all months except April in southwestern São Paulo state, Brazil; lactating specimens appeared from November to May. A protracted reproductive season and polyestry, indicated by simultaneously pregnant and lactating females in November and December, characterized the São Paulo population. Willig (1985) defined reproductive patterns in two well-delineated populations from northeast Brazil. Differences in the annual reproductive cycle between Caatingas and edaphic Cerrado populations were not evident. A bimodal distribution of pregnancy began in the early dry season (July) and terminated early in the rainy season (February–March). Pregnancy was no longer observed by the end of the rainy season. A bimodal distribution of lactation also was evident; lactating females were first observed at the end of the dry season (October) and continued to be captured into the middle of the rainy season.

FORM AND FUNCTION. Specimens from Bahia were reported by McNab and Morrison (1963) to contain relatively large salivary glands (1.3 to 2.5% of body mass) compared to other phyllostomids (for example, *Chiroderma villosus*, 0.6% of body mass). McNab (1982) reported that *V. lineatus* maintains body temperature independently of ambient temperatures to at least 10°C, with body temperature elevating in response to increases in environmental temperature beyond 30°C.

ECOLOGY AND BEHAVIOR. Based upon the works of Ruschi (1953) and McNab (1969), Gardner (1977a) defined the diet of this species as including fruits and insects, especially sphingid moths. McNab and Morrison (1963) occasionally captured *V. lineatus* near sapotes (*Achras zapota*) but did not report if the fruit was actually consumed. Taddei (1973) reported that *Carica papaya*, *Cecropia*, *Chlorophora tinctoria*, *Ficus enormis*, *Ficus* sp., and *Syzygium jambos* composed the natural diet in northwestern São Paulo state, Brazil; banana and honey water were consumed in captivity. McNab and Morrison (1963) previously reported a preference in captivity for papaya over banana. Willig (1983) suggested that *Vismia* was an important dietary component on the Chapada do Araripe in Ceará state, Brazil, because it was the dominant food item in stomachs and feces examined. The observations of Sazima and Sazima (1975) and Sazima (1976) expanded the reported feeding niche of *V. lineatus* to include nectarivory; *Musa acuminata* and *Lafroesia pacari* flowers often were exploited by *V. lineatus* on Serra do Cipó, in the Brazilian state of Minas Gerais. Sazima and Sazima (1975) further detailed the feeding behavior of the species. Specimens arrived at flowering *Lafroesia* at dusk, made one or two short passes and inserted their heads, while in flight, into the corolla. When the flower was inclined horizontally, *V. lineatus* suspended itself by the thumbs alone from the base of the flower. When the flower was vertical, *V. lineatus* also supported itself by anchoring the feet on the branch supporting the inflorescence. A sequence of rapid head movements followed, during which time individuals lapped nectar and pollen. Each visit lasted from 0.5 to 1.5 s. Stages of floral development previous to anthesis were exploited also. Group and solitary foraging was observed, the latter more common on plants with few available flowers.

Willig (1983) reported unequal sex ratios for populations in northeast Brazil. Moreover, he suggested that male *V. lineatus* maintained small harems of 7 to 15 females in small granitic caves. Similar roosting colonies from 6 to approximately 20 individuals also were reported by Peracchi and Albuquerque (1971). They observed the smaller groups roosting under leaves of the palm, *Livistona oliviformis*, and larger groups roosting under *Syagrus picrophylla* leaves. Taddei (1973) observed eight individuals roosting beneath a *Dracaena* leaf (approximately 3 mm in height) and two individuals in a bromeliad 12 m off the ground.

Vampyrops lineatus has been captured with *Anoura geoffroyi*, *Artibeus concolor*, *A. fimbriatus*, *A. jamaicensis*, *A. lituratus*, *Carollia perspicillata*, *Chiroderma doriae*, *C. vellosus*, *Chrotopterus auritus*, *Desmodus rotundus*, *Diphylla ecaudata*, *Eptesicus brasiliensis*, *E. diminutus*, *E. furinalis*, *Eumops auri- pendulus*, *E. bonariensis*, *E. glaucinus*, *E. perotis*, *Furipterus horrens*, *Glossophaga soricina*, *Histiotus velatus*, *Lasiurus borealis*, *L. cinereus*, *L. ega*, *Lonchophylla mordax*, *Micronycteris megalotis*, *M. minuta*, *Mimon crenulatum*, *Molossops abrasus*, *M. planirostris*, *M. temminckii*, *Molossus ater*, *M. molossus*, *Myotis albecens*, *M. nigricans*, *M. riparius*, *M. ruber*, *M. simus*, *Natalus stramineus*, *Neoplatymops mattogrossensis*, *Noctilio al-*



FIG. 4. A single diploid chromosomal complement of a male *V. l. lineatus* (Univ. Michigan number 125428); the sex chromosomes are in the lower right corner.

biventris, *N. leporinus*, *Peropteryx macrotis*, *Phyllostomus discolor*, *P. hastatus*, *Promops nasutus*, *P. occultus*, *Pteronotus davyi*, *Pygoderma bilabiatum*, *Saccopteryx leptura*, *Sturnia lilium*, *Tadarida brasiliensis*, *T. laticaudata*, *Tonatia bidens*, *T. brasiliense*, *T. silvicola*, *Trachops cirrhosus*, *Uroderma magnirostrum*, and *Vampyressa pusilla* (Myers, 1982; Taddei, 1973; Willig, 1983).

GENETICS. Figure 4 illustrates a single diploid chromosomal complement from a male *V. l. lineatus* from Paraguay (courtesy of P. Myers). The chromosomal morphology is identical to that reported for *V. l. nigellus* (as *V. nigellus* by Gardner, 1977b) and all other congeners (Baker, 1979; Baker et al., 1982). The only variation in chromosomal morphology that has been documented for the genus *Vampyrops* is that the Y chromosome of some species (*V. brachycephalus*, *V. dorsalis*, and *V. helleri*) is submetacentric, whereas in other species (*V. infuscus*, *V. l. nigellus*, and *V. vittatus*), it is acrocentric (Baker, 1979; Baker et al., 1982; Gardner, 1977b). The condition of the Y chromosome of *V. l. lineatus* is acrocentric like that of *V. l. nigellus* reported by Gardner (1977b). The karyotype of *V. lineatus* and all reported congeners consists of a diploid number of 30 with a fundamental number of 56. Electrophoretic analyses involving *V. lineatus* have not been reported.

REMARKS. The English vernacular name, Geoffroy's Rayed Bat, is derived from a literal translation of the French vulgar name, "Phyllostome rayé," written on the wing of the type specimen by Geoffroy. The generic name *Vampyrops* is derived from the combination of the Russian root "vampyr" meaning a vampire and the Greek root "ops" meaning appearance. The specific epithet is derived from the Latin root "linea" meaning a line. Thus, the scientific name *Vampyrops lineatus* means a vampire-appearing bat with a line.

Foremost, we would like to acknowledge J. K. Jones, Jr., for his advice and taxonomic expertise. C. Jones, C. Hood, P. Myers, M. B. Qumseyeh, and D. Wilson provided critical reviews of the manuscript. In addition, P. Myers provided the slides with chromosomal preparations for us to photograph; his kindness is appreciated. J. Roach expertly typed the manuscript.

LITERATURE CITED

- ALLEN, H. 1891. Description of a new species of *Vampyrops*. Proc. Acad. Nat. Sci. Philadelphia, 43:400–405.
- ANDERSON, S., K. F. KOOPMAN, AND G. K. CREIGHTON. 1982. Bats of Bolivia: an annotated checklist. Amer. Mus. Novitates, 2750:1–24.
- BAKER, R. J. 1979. Karyology. Pp. 107–155, in Biology of bats of the New World family Phyllostomatidae. Part III (R. J. Baker, J. K. Jones, Jr., and D. C. Carter, eds.). Spec. Publ. Mus., Texas Tech Univ., 16:1–441.
- BAKER, R. J., M. W. HAIDUK, L. W. ROBBINS, A. CADENA, AND B. F. KOOP. 1982. Chromosomal studies of South American bats and their systematic implications. Pp. 303–327, in Mam-

- malian biology in South America (M. A. Mares and H. H. Genoways, eds.). Spec. Publ. Ser., Pymatuning Lab. Ecol., Univ. Pittsburgh, 6:1-539.
- CABRERA, A. 1958. Catálogo de los mamíferos de America del Sur. Rev. Mus. Argentino Cien. Nat. "Bernardino Rivadavia," Cien. Zool., 4:1-307.
- CARTER, D. C., AND P. G. DOLAN. 1978. Catalogue of type specimens of Neotropical bats in selected European museums. Spec. Publ. Mus., Texas Tech Univ., 15:1-136.
- CHUNA VIEIRA, C. O. DA. 1942. Ensaio monográfico sobre os Quirópteros do Brasil. Arquivos Zool., Estado de São Paulo, 3:219-471.
- DOBSON, G. E. 1878. Catalogue of the Chiroptera in the collection of the British Museum. British Mus., London, 567 pp.
- ELLIOT, D. G. 1904. The land and sea mammals of Middle America and the West Indies. Field Columbian Mus., Zool. Ser., 4: 441-850.
- GARDNER, A. L. 1977a. Feeding habits. Pp. 293-350, in Biology of bats of the New World family Phyllostomatidae. Part II (R. J. Baker, J. K. Jones, Jr., and D. C. Carter, eds.). Spec. Publ. Mus., Texas Tech Univ., 13:1-364.
- . 1977b. Chromosomal variation in *Vampyressa* and a review of chromosomal evolution in the Phyllostomidae (Chiroptera). Syst. Zool., 26:300-318.
- GARDNER, A. L., AND D. C. CARTER. 1972. A new stenodermine bat (Phyllostomatidae) from Peru. Occas. Papers Mus., Texas Tech Univ., 2:1-4.
- GEOFFROY ST.-HILAIRE, É. 1810. Sur les Phyllostomes et les Megadermes, deux genres de la famille des chauve-souris. Ann. Mus. Nat. Hist. Nat., Paris, 15:157-198 + pls. ix-xii.
- GOODWIN, G. G. 1946. Mammals of Costa Rica. Bull. Amer. Mus. Nat. Hist., 87:271-471.
- HERSHKOVITZ, P. 1949. Mammals of northern Colombia. Preliminary report no. 5: bats (Chiroptera). Proc. U.S. Natl. Mus., 99:429-454.
- HONACKI, J. H., K. E. KINMAN, AND J. W. KOEPL (EDS.). 1982. Mammal species of the World: a taxonomic and geographic reference. Allen Press, Inc., and The Assoc. Syst. Coll., Lawrence, Kansas, 694 pp.
- JONES, J. K., JR., AND D. C. CARTER. 1976. Annotated checklist with keys to subfamilies and genera. Pp. 7-38, in Biology of bats of the New World family Phyllostomatidae. Part I (R. J. Baker, J. K. Jones, Jr., and D. C. Carter, eds.). Spec. Publ. Mus., Texas Tech Univ., 10:1-218.
- . 1979. Systematic and distributional notes. Pp. 7-11, in Biology of bats of the New World family Phyllostomatidae. Part III (R. J. Baker, J. K. Jones, Jr., and D. C. Carter, eds.). Spec. Publ. Mus., Texas Tech Univ., 16:1-441.
- KOOPMAN, K. F. 1978. Zoogeography of Peruvian bats with special emphasis on the role of the Andes. Amer. Mus. Novitates, 2651:1-33.
- . 1982. Biogeography of the bats of South America. Pp. 273-302, in Mammalian biology in South America (M. A. Mares and H. H. Genoways, eds.). Spec. Publ. Ser., Pymatuning Lab. Ecol., Univ. Pittsburgh, 6:1-539.
- LIMA, J. O. 1926. Os morcegos da coleção do Museu Paulista. Rev. Mus. Paulista, São Paulo, 14:43-127.
- MCNAB, B. K. 1969. The economics of temperature regulation in Neotropical bats. Comp. Biochem. Physiol., 31:227-268.
- . 1982. Evolutionary alternatives in the physiological ecology of bats. Pp. 151-200, in Ecology of bats (T. H. Kunz, ed.). Plenum Press, New York, 425 pp.
- MCNAB, B. K., AND P. MORRISON. 1963. Observations on bats from Bahia, Brazil. J. Mamm., 44:21-23.
- MYERS, P. 1982. Origins and affinities of the mammal fauna of Paraguay. Pp. 85-93, in Mammalian biology in South America (M. A. Mares and H. H. Genoways, eds.). Spec. Publ. Ser., Pymatuning Lab. Ecol., Univ. Pittsburgh, 6:1-539.
- PERACCHI, A. L., AND S. T. DE ALBUQUERQUE. 1971. Lista provisória dos quirópteros dos estados do Rio de Janeiro e Guanabara, Brasil (Mammalia, Chiroptera). Rev. Brasil. Biol., 31: 405-413.
- PETERS, W. 1865. Monatsberichte Koniglichen Preufs. Akademie de Wissen., Berlin. 1865, p. 356.
- ROUK, C. S., AND D. C. CARTER. 1972. A new species of *Vampyrops* (Chiroptera: Phyllostomatidae) from South America. Occas. Papers Mus., Texas Tech Univ., 1:1-7.
- RUSCHI, A. 1953. Morcegos do estado do Espirito Santo. XVIII. Familia Phyllostomidae. Descrição das especies *Artibeus jamaicensis lituratus* e *Vampyrops lineatus*, com algumas observações. Bol. Mus. Biol. Prof. Mello-Leitão, Santa Teresa, Zool., 20:1-8.
- SANBORN, C. C. 1955. Remarks on the bats of the genus *Vampyrops*. Fieldiana Zool., 37:403-413.
- SAZIMA, I. 1976. Observations on the feeding habits of phyllostomid bats (*Carollia*, *Anoura*, and *Vampyrops*) in southeastern Brazil. J. Mamm., 57:381-382.
- SAZIMA, M., AND I. SAZIMA. 1975. Quiróptero-filia em *Lafoensia pacari* St. Hil. (Lythraceae), na Serra do Cipó, Minas Gerais. Ciência e Cultura, 27:406-416.
- SWANEPOEL, P., AND H. H. GENOWAYS. 1979. Morphometrics. Pp. 13-106, in Biology of bats of the New World family Phyllostomatidae. Part III (R. J. Baker, J. K. Jones, Jr., and D. C. Carter, eds.). Spec. Publ. Mus., Texas Tech Univ., 16: 1-441.
- TADDEI, V. A. 1973. Phyllostomidae da Região Norte-ocidental do Estado de São Paulo. Unpubl. Ph.D. dissert., Univ. Estadual Paulista de São José do Rio Preto, São Paulo, 249 pp.
- . 1976. The reproduction of some Phyllostomidae (Chiroptera) from the northwestern region of the State of São Paulo. Bol. Mus. Zool., Univ. São Paulo, 1:313-330.
- . 1979. Phyllostomidae (Chiroptera) do Norteocidental do Estado de São Paulo III—Stenodermatinae. Ciência e Cultura, 31:900-914.
- WILLIG, M. R. 1983. Composition, microgeographic variation, and sexual dimorphism in Caatingas and Cerrado bat communities from Northeast Brazil. Bull. Carnegie Mus. Nat. Hist., 23:1-131.
- . 1985. Reproductive patterns of bats from Caatingas and Cerrado biomes of Northeast Brazil. J. Mamm., 66:668-681.

Editors of this account were B. J. VERTS and J. K. JONES, JR. Managing editor was CARLETON J. PHILLIPS.

M. R. WILLIG AND R. R. HOLLANDER, DEPARTMENT OF BIOLOGICAL SCIENCES AND THE MUSEUM, TEXAS TECH UNIVERSITY, LUBBOCK 79409.